

Free Groups in Quaternion Algebras

S. O. Juriaans and A. C. Souza Filho

ABSTRACT. In [6] we constructed pairs of units u, v in a $\mathbb{Z}$ -order of a quaternion algebra over imaginary rational extensions $\mathbb{Q}\sqrt{-d}$, where $d \equiv 7 \pmod{8}$ is positive and square free integer, such that the group generated by u^n, v^n is a free group for some positive integer n . Here we extend this result to any imaginary quadratic extension of $\mathbb{Q}$, thus including matrix algebras. We show that $\langle u^n, v^n \rangle$ is a free group for all non-zero integers n , except when $d = 2$ yielding $n > 1$. The units we use in our construction belong to a new class of units, coming from Pell's and Gauss' equations. Also, we establish a criterium for a pair of elements to generate a free semigroup.

INTRODUCTION

Constructions of free groups is a nontrivial problem and is a main research topic. Let R be a ring with unity and G a group, denote by RG the set $\{r_1g_1 + \cdots + r_ng_n, r_i \in R, g_i \in G, 0 < i \leq n, n \in \mathbb{N}\}$ of formal sums in RG . Clearly, RG is a unitary ring. Denote by $\mathcal{U}(RG)$ the group of units of the ring RG and $\mathcal{U}_1(RG)$ the subset of $\mathcal{U}(RG)$, such that, if $u = u_1g_1 + \cdots + u_ng_n$ is in $\mathcal{U}_1(RG)$, then $u_1 + \cdots + u_n = 1$. The set $\{u_1, \dots, u_n\} =: \text{supp}(u)$ is the support of $u \in \mathcal{U}(RG)$. In [5], Higman shows that if A is a finite abelian group, then $\mathcal{U}_1(\mathbb{Z}A) = A \times F$, where F is a finitely generated free abelian group. In [4], it is proved that if G is a finite non-Hamiltonian 2-group, then $\mathcal{U}(\mathbb{Z}G)$ contains a free group. On the other hand, a result of Tits, [11], gives necessary conditions for a pair of diagonalizable elements of $GL(2, \mathbb{C})$ to generate a free group. In [9], assuming the existence of nontrivial units in $\mathbb{Z}G$ defined by bicyclic units, free groups are constructed in $\mathcal{U}(\mathbb{Z}G)$. In [7], pairs of units are constructed generating a free group in quaternion algebras. In [8], the same was done in finite dimensional crossed products.

In [6], finite groups G and rings R of characteristic zero, with $\mathcal{U}(RG)$ a hyperbolic group, were classified. Units, arising from solutions of Pell's and Gauss' equations, were also constructed in the algebras $(\frac{-1, -1}{R})$, where $R = \mathfrak{o}_{\mathbb{Q}(\sqrt{-d})}$ is the ring of algebraic integers of quadratic extensions of $\mathbb{Q}$. We coined the names Pell and Gauss units. In [6], we also use a Gauss unit of norm -1 and the deep results of [3], to give a full set of generators of the unit group of $\mathbb{Z}[\frac{1+\sqrt{-7}}{2}]K_8$, where K_8 is the quaternion group of order 8. Using a result of Gromov, we show that this unit group

1991 *Mathematics Subject Classification.* Primary 16U60, 20E05; Secondary 16S34, 20F67.

Key words and phrases. Hyperbolic Groups, Quaternion Algebras, Free Groups, Free Semigroups, Group Rings, Units, Möbius transformation.

The first author is supported by CNPq and the second author by FAPESP-Brazil, Proc. 2008/57930-1.

is a hyperbolic group with one end and whose hyperbolic boundary is the 2-dimensional euclidean sphere. From this and [2, Proposition III.Γ.3.20], it follows that if $d \equiv 7 \pmod{8}$ is a positive integer, and $u, v \in \mathcal{U}((\frac{-1, -1}{R}))$ are Pell units with distinct supports then $\langle u^n, v^n \rangle$ is a free group for a suitable positive n .

The problem on determining the power n , is a non-trivial task. In general, as in [2, Proposition III.Γ.3.20], we have results about the existence of the power n . In the section 2, we construct units with suitable algebraic properties. With a pair of these units, we can use the Ping-Pong Lemma, in a non-standart way, to determine the power n , hence generalizing [6, Theorem 5.5].

Here, we construct free groups of rank 2 in $\mathbb{Z}$ -orders of quaternion algebras over all imaginary quadratic extensions of the rational number field, in particular those which are division rings. The pair of 2-Pell units we use is such that we can neither apply Proposition 3.20 of [2] nor Tits' result because the norm of the eigenvalues of the units, when considered as matrices, is equal to one. We finish the paper with a result which gives a criterium for two elements to generate a free semigroup. We show how it is applied in our case.

1. Preliminaries

We denote by $(\frac{-1, -1}{K}) := K[i, j : i^2 = -1, j^2 = -1, -ji = ij =: k]$ the quaternion algebra over K . Let K be an algebraic number field and $\mathfrak{o}_K$ its ring of integers, denote by $(\frac{-1, -1}{\mathfrak{o}_K})$, the $\mathfrak{o}_K$ -span of $\{1, i, j, k\}$, which is an $\mathfrak{o}_K$ -algebra. Clearly, since $\mathfrak{o}_K$ is a $\mathbb{Z}$ -order of K , the algebra $(\frac{-1, -1}{\mathfrak{o}_K})$ is a $\mathbb{Z}$ -order of $(\frac{-1, -1}{K})$.

We define the monomorphism σ over $\mathcal{H} := (\frac{-1, -1}{\mathbb{C}})$, $\sigma : \mathcal{H} \rightarrow \mathcal{H}$, $\sigma(x) = j^{-1}xj$.

Clearly, the quaternion algebra $\mathcal{H}$ is isomorphic to $\mathbb{C} \otimes \mathbb{C}j$, where $i \in \mathcal{H}$ is associated to $\sqrt{-1} \in \mathbb{C}$ and $\otimes$ denotes a cross product over $\mathbb{C}$. Since, all elements $z \in \mathcal{H}$ is of the form $z = x + yj$, where $x, y \in \mathbb{C} + \mathbb{C}[i]$, the $\mathbb{C}$ -monomorphism $\Psi : \mathcal{H} \hookrightarrow M_2(\mathbb{C})$, $\Psi(x + yj) = \begin{pmatrix} x & y \\ -\sigma(y) & \sigma(x) \end{pmatrix}$ is well defined. We may identify $\sigma(x) = \sigma(a + bi) = a - bi$ with the complex number $a - b\sqrt{-1}$. Thus, a quaternion algebra is isomorphic to a complex matrix of the form $\begin{pmatrix} x & y \\ -\bar{y} & \bar{x} \end{pmatrix}$, where $\bar{x}$ denotes the conjugation of x induced by σ , see also [1].

If $M = (m_{ij}) \in GL_2(\mathbb{C})$, then M defines a Möbius transformation $\varphi_M(z) := \frac{m_{11}z + m_{12}}{m_{21}z + m_{22}}$. Thus Ψ induces a homomorphism $\varphi_u : \mathbb{C} \hookrightarrow \mathcal{M}$, $\varphi_u := \varphi_{\Psi(u)}$ and $\mathcal{M}$ is the group of Möbius transformations.

We restrict the maps σ and Ψ on the quaternion algebra $(\frac{-1, -1}{\mathbb{Q}\sqrt{-d}}) \subset \mathcal{H}$. Hence $\Psi((\frac{-1, -1}{\mathbb{Q}\sqrt{-d}})) \subset M_2(K)$, where $K = \mathbb{Q}(\sqrt{-1}, \sqrt{-d})$ the $\mathbb{Q}$ -span of $\{1, \sqrt{-1}, \sqrt{-d}, \sqrt{d}\}$. If $u = u_1 + u_i i + u_j j + u_k k \in (\frac{-1, -1}{\mathbb{Q}(\sqrt{-d})})$, then $\Psi((u_1 + u_i i) + (u_j + u_k i)j) = \begin{pmatrix} u_1 + u_i \sqrt{-1} & u_j + u_k \sqrt{-1} \\ -u_j + u_k \sqrt{-1} & u_1 - u_i \sqrt{-1} \end{pmatrix}$, defines the Möbius transformations $\varphi_u(z)$, where $z \in \mathbb{C}$.

In [10] it is shown that $\langle z + 2, \frac{z}{2z+1} \rangle$ is a free group of the group $\mathcal{M}$. This is a trivial application of the Ping-Pong Lemma, [2, Lemma 3.19], which states that if $h_1, \dots, h_r$ are bijections of a set Ω and there exist non-empty disjoint subsets $A_{1,1}, A_{1,-1}, \dots, A_{r,1}, A_{r,-1} \subset \Omega$ such that $h_i^\epsilon(\Omega \setminus A_{i,\epsilon}) \subset A_{i,-\epsilon}$, for $\epsilon \in \{-1, 1\}$ and $i = 1, \dots, r$, then $\langle h_1, \dots, h_r \rangle$ generates a free subgroup of rank r in $\text{Perm}(\Omega)$. In fact, $h_1(z) := \frac{z}{2z+1}$ and $h_2(z) := z + 2$ are bijections of $\Omega := \mathbb{R} \cup \{-\infty, \infty\}$. It is easily verified that the sets $A_{1,1} :=]-1, 0[$, $A_{1,-1} := [0, 1]$, $A_{2,1} := [-\infty, -1]$, and $A_{2,-1} :=]1, \infty[$ are in the conditions of the Ping-Pong Lemma.

Theorem 1.1. *The group generated by the units $u = \sqrt{-1} + (-\sqrt{-1} + i)j$ and $w = \sqrt{-1} + (\sqrt{-1} + i)j$ in $(\frac{-1, -1}{\mathfrak{o}_{\mathbb{Q}(\sqrt{-1})}})$ is free.*

PROOF. Consider the images of u and w in $M_2(\mathbb{C})$ which are $\Psi(u) = \begin{pmatrix} \sqrt{-1} & 0 \\ 2\sqrt{-1} & \sqrt{-1} \end{pmatrix}$ and $\Psi(w) = \begin{pmatrix} \sqrt{-1} & 2\sqrt{-1} \\ 0 & \sqrt{-1} \end{pmatrix}$. As Möbius transformations they generate the group $\langle \frac{z}{2z+1}, z+2 \rangle$ which is free of rank two. $\square$

2. Free Groups in Quaternion Algebras

In the sequel, $K = \mathbb{Q}(\sqrt{-d})$ is an imaginary quadratic extension with d positive and square-free integer. Let $\xi \neq \psi$ be elements of $\{1, i, j, k\}$. Suppose

$$(2.1) \quad u := m\sqrt{-d}\xi + p\psi, \text{ where } p, m \in \mathbb{Z},$$

is an element in $(\frac{-1, -1}{\mathfrak{o}_K})$ having norm 1. Then

$$(2.2) \quad p^2 - m^2d = 1,$$

i.e., the pair (p, m) is a solution of Pell's equation $x^2 - dy^2 = 1$ in $\mathbb{Q}(\sqrt{d})$. Equation (2.2) implies that $\epsilon = p + m\sqrt{d}$ is a unit in $\mathfrak{o}_{\mathbb{Q}(\sqrt{d})}$. Conversely, if $\epsilon = x + y\sqrt{d}$ is a unit of norm 1 in $\mathfrak{o}_{\mathbb{Q}(\sqrt{d})}$ then, necessarily, $x^2 - y^2d = 1$, and, therefore, for any choice of ξ, ψ in $\{1, i, j, k\}$, $\xi \neq \psi$,

$$(2.3) \quad y\sqrt{-d}\xi + x\psi$$

is a unit in $(\frac{-1, -1}{\mathfrak{o}_K})$. In particular,

$$(2.4) \quad u_{(\epsilon, \psi)} := x + y\sqrt{-d}\psi, \quad \psi \in \{i, j, k\}.$$

Thus $u_{(\epsilon, \psi)}$ is a unit in $(\frac{-1, -1}{\mathfrak{o}_K})$.

With the notations as above, we have:

Proposition 2.1. [6]

- (1) If $1 \notin \text{supp}(u)$, the support of u , then u is a torsion unit.
- (2) If $\epsilon = x + y\sqrt{d}$ is a unit in $\mathfrak{o}_{\mathbb{Q}(\sqrt{d})}$, then

$$u_{(\epsilon, \psi)}^n = u_{(\epsilon^n, \psi)}$$

for all $\psi \in \{i, j, k\}$ and $n \in \mathbb{Z}$.

Units of type (2.3) are called *2-Pell units*. Denote the norm of $\epsilon = x + y\sqrt{d}$ by $\mathcal{N}(\epsilon) := \epsilon\bar{\epsilon}$, where $\bar{\epsilon} = x - y\sqrt{d}$. If $\mathcal{N}(\epsilon) = -1$, then the unit $y\sqrt{-d}\xi + x\psi$ is also a 2-Gauss unit, see [6].

Likewise, for $\mathcal{N}(\epsilon) = 1$, we define the 4-Pell unit $u := \frac{y}{2}\sqrt{-d}\zeta + (\frac{y}{2}\sqrt{-d})\xi + (\frac{1+x}{2})\psi + (\frac{1-x}{2})\phi$, where $\zeta, \xi, \psi, \phi \in \{1, i, j, k\}$ are different of each other.

Remark 2.2.

- We cannot define a 4-Pell unit when $\mathcal{N}(\epsilon) = -1$, since the unit has norm $\frac{2+2(x^2-y^2d)}{4} \neq 0$.

- When $y \equiv 1 \pmod{2}$, the invertible ϵ^2 defines a 4-Pell unit $u := xy\sqrt{-d} + (xy\sqrt{-d})i + (x^2)j + (y^2d)k$.
- If the invertible $\epsilon = (2x - 1) + y\sqrt{2d} \in \mathbb{Q}(\sqrt{2d})$ has norm $\mathcal{N}(\epsilon) = 1$, we define a 3-Pell unit $u := y\sqrt{-d}\xi + x\psi + (1 - x)\phi$ in $(\frac{-1, -1}{\sigma_{\mathbb{Q}(\sqrt{-d})}})$, where $\xi, \psi, \phi \in \{1, i, j, k\}$ are different of each other.

Let $\epsilon = x + y\sqrt{d}$ be the fundamental invertible, $u = x + (y\sqrt{-d})i$ and $\varphi_u(z) = \frac{x-y\sqrt{d}}{x+y\sqrt{d}}z$ its image in $\mathcal{M}$. Since x, y are always positive, if $\mathcal{N}(\epsilon) = 1$, then $x + y\sqrt{d} > 1$ and $0 < x - y\sqrt{d} < 1$. Hence, we have that $\varphi_u(z) = \rho z$ with $\rho = \frac{x-y\sqrt{d}}{x+y\sqrt{d}} \in]0, 1[$.

Let $\varphi(z) = \frac{az+b}{cz+d}$. Define $z_p := \frac{-d}{c}$, respectively, $z_0 := \frac{-b}{a}$, the pole and the zero of φ . Given a Pell unit w , the points $(z_0, 0)$, (z_p, φ_∞) , when $\varphi \in \{\varphi_w, \varphi_w^{-1}\}$, play an important role in what follows.

For suitable positive real numbers m_2, m_1, n_1, n_2 , a disjoint partition of the extended real numbers is either:

$[-\infty, m_2 z_p[\dot{\cup} [m_2 z_p, m_1 z_0[\dot{\cup} [m_1 z_0, n_1 z'_0[\dot{\cup} [n_1 z'_0, n_2 z'_p[\dot{\cup} [n_2 z'_p, \infty]$ if $z_p < z_0 < 0 < z'_0 < z'_p$, or:
 $[-\infty, m_2 z_0[\dot{\cup} [m_2 z_0, m_1 z_p[\dot{\cup} [m_1 z_p, n_1 z'_p[\dot{\cup} [n_1 z'_p, n_2 z'_0[\dot{\cup} [n_2 z'_0, \infty]$ if $z_0 < z_p < 0 < z'_p < z'_0$, where z_0 is the zero and z_p is the pole of φ_w , and z'_0 is the zero and z'_p is the pole of φ_w^{-1} . Our next result proves, on the conditons of the theorem, that φ_u , φ_w and the sets above satisfy the Ping-Pong Lemma. Since φ is a continuous mapping with positive derivative, if $a_2 < z_0 < z_p < a_1 < 0$, $\frac{a}{c} > 0$ and $A_{11} :=]a_2, a_1[$, then $\varphi(\mathbb{R} \setminus A_{11}) =]\varphi(a_2), \varphi(a_1)[$. In the proof of the next theorem we freely use this property.

Theorem 2.3. *Let $\epsilon = x + y\sqrt{d}$ be the fundamental invertible in $\mathbb{Q}(\sqrt{d})$ of norm $\mathcal{N}(\epsilon) = \epsilon\bar{\epsilon} = 1$ and $u = x + (y\sqrt{-d})i$. If w is one of the units*

- (1.): $y\sqrt{-d} + (x)k$;
- (2.): $\frac{x+1}{2} - (\frac{y\sqrt{-d}}{2})i + (\frac{x-1}{2})j + (\frac{y\sqrt{-d}}{2})k$;
- (3.): $x^2 - (xy\sqrt{-d})i - (y^2d)j + (xy\sqrt{-d})k$.

Then the group $\langle u, w \rangle \subset \mathcal{U}((\frac{-1, -1}{\sigma_K}))$ is a free group.

PROOF. Consider $\varphi_u, \varphi_w \in \mathcal{M}$ and $\Omega := \mathbb{R} \cup \{\pm\infty\}$. We claim that there exist real numbers $a_2 < a_1 < 0 < b_1 < b_2$ such that

$$A_{1,1} := [a_2, a_1], \quad A_{1,-1} := [b_1, b_2], \quad A_{2,1} := [-\infty, a_2[\dot{\cup}]b_2, \infty] \quad \text{and} \quad A_{2,-1} :=]a_1, b_1[$$

are sets satisfying the conditions stated in the Ping-Pong Lemma.

In fact, let $h_2(z) := \varphi_u(z) = \frac{x-y\sqrt{d}}{x+y\sqrt{d}}z$ and $h_1(z) := \varphi_w(z)$, where w is like in the itens 1, 2, 3. Then since each $A_{i,\pm 1}$, $i = 1, 2$ is to be an interval, or a disjoint union of two intervals, we get the following conditions.

- first:** $h_1(\Omega \setminus A_{1,1}) \subset A_{1,-1}$, which is equivalent to $h_1(a_2), h_1(a_1) \in A_{1,-1}$;
- second:** $h_1^{-1}(\Omega \setminus A_{1,-1}) \subset A_{1,1}$, which is equivalent to $h_1^{-1}(b_1), h_1^{-1}(b_2) \in A_{1,1}$;
- third:** $h_2(\Omega \setminus A_{2,1}) \subset A_{2,-1}$, which is equivalent to $h_2(a_2), h_2(b_2) \in A_{2,-1}$;
- fourth:** $h_2^{-1}(\Omega \setminus A_{2,-1}) \subset A_{2,1}$, which is equivalent to $h_2^{-1}(a_1), h_2^{-1}(b_1) \in A_{2,1}$.

(1) Set $h_1(z) := \frac{x}{y\sqrt{d}} \frac{\frac{y\sqrt{d}}{x}z+1}{\frac{x}{y\sqrt{d}}z+1}$, $-a_2 = b_2 := \frac{3}{2} \frac{x}{y\sqrt{d}}$ and $-a_1 = b_1 := \frac{1}{2} \frac{y\sqrt{d}}{x}$. The first condition: $h_1(\Omega \setminus A_{1,1}) =]h_1(a_2), h_1(a_1)[$; we calculate $h_1(a_2) = \frac{xy\sqrt{d}}{x^2+2}$ and $h_1(a_1) = \frac{x^2+1}{xy\sqrt{d}}$. Since $x^2 - y^2d = 1$,

we have $b_1 < h_1(a_2)$. Similarly, we compute $h_1(a_1) = \frac{x^2+1}{xy\sqrt{d}} < \frac{3x}{2y\sqrt{d}} = b_2$, thus $h_1(\Omega \setminus A_{1,1}) \subset A_{1,-1}$. The second condition: $h_1^{-1}(\Omega \setminus A_{1,-1}) =]h_1^{-1}(b_2), h_1^{-1}(b_1)[$; we compute $h_1^{-1}(b_2) = \frac{-xy\sqrt{d}}{x^2+2} = -h_1(a_2)$ and $h_1^{-1}(b_1) = \frac{-x^2-1}{xy\sqrt{d}}$. Clearly, $h_1^{-1}(b_2) < a_1$ and $a_2 < h_1^{-1}(b_1)$, thus $h_1^{-1}(\Omega \setminus A_{1,-1}) \subset A_{1,1}$. The third and fourth conditions: $h_2(\Omega \setminus A_{2,1}) = [h_2(a_2), h_2(b_2)]$ and clearly $h_2(b_2) = -h_2(a_2)$. Denoting by $()$ the order relation between $h_2(b_2) = \frac{3x}{2y\sqrt{d}}\rho$ and b_1 we have that $\frac{3x}{2y\sqrt{d}}\frac{x-y\sqrt{d}}{x+y\sqrt{d}}() \frac{y\sqrt{d}}{2x}$. Since the real numbers $h_2(b_2)$ and b_1 are positive, it follows that $\frac{3x^2}{y^2d}() \frac{x+y\sqrt{d}}{x-y\sqrt{d}}$. Clearly, $\frac{3x^2}{y^2d} = 3 + \frac{3}{y^2d} < 5 < x^2 + y^2d + 2xy\sqrt{d} = \frac{x+y\sqrt{d}}{x-y\sqrt{d}}$. Thus $h_2(b_2) < b_1$ and $a_1 < h_2(a_2)$. Since $h_2^{-1}(z) = \frac{1}{\rho}z$ we proved that $h_2(\Omega \setminus A_{2,1}) \subset A_{2,-1}$ and $h_2^{-1}(\Omega \setminus A_{2,-1}) = [-\infty, h_2^{-1}(a_1)] \cup [h_2^{-1}(b_1), \infty] \subset A_{2,1}$.

(2) In this case the pole and root of $h_1(z) = \frac{(y\sqrt{d}+(x+1))z-(y\sqrt{d}-(x-1))}{-(y\sqrt{d}+(x-1))z-(y\sqrt{d}-(x+1))}$ are, respectively $z_p = \frac{-(y\sqrt{d}-(x+1))}{y\sqrt{d}+(x-1)}$ and $z_0 = \frac{y\sqrt{d}-(x-1)}{y\sqrt{d}+(x+1)}$ and the pole and the root of φ_w^{-1} are $z'_p = \frac{-(y\sqrt{d}+(x+1))}{y\sqrt{d}+(x-1)}$ and $z'_0 = \frac{y\sqrt{d}-(x-1)}{y\sqrt{d}-(x+1)}$. Since z_p is positive, define the intervals by $A_{1,-1} = [a_2, a_1]$ and $A_{1,1} := [b_1, b_2]$

with $a_2 := 3z'_p$, $a_1 := \frac{z'_0}{2}$ and $b_1 := \frac{z_0}{2}$, $b_2 := 3z_p$, and proceed as before proving that $h_1(\Omega \setminus A_{1,1}) =]h_1(b_2), h_1(b_1)[\subset A_{1,-1}$ and $h_1^{-1}(\Omega \setminus A_{1,-1}) =]h_1^{-1}(a_1), h_1^{-1}(a_2)[\subset A_{1,1}$. This is true, provide that $\frac{x}{y\sqrt{d}} < \frac{7x+5}{5x+7}$, if $x > 2$. This holds because the congruence $y \equiv 0 \pmod{2}$ implies that $x > 2$.

(3) We have that $h_1(z) = \frac{(x+y\sqrt{d})}{x-y\sqrt{d}} \frac{xz-y\sqrt{d}}{-(y\sqrt{d})z+x}$. Set $b_1 := \frac{z_0}{2} = \frac{y\sqrt{d}}{2x}$, $b_2 := 3z_p = \frac{x}{y\sqrt{d}}$, $a_1 := \frac{z'_0}{2} = -\frac{b_1}{\rho}$ and $a_2 := 3z'_p = -\frac{b_2}{\rho}$, where $\rho = \frac{x-y\sqrt{d}}{x+y\sqrt{d}}$ and the intervals $A_{1,-1} := [a_2, a_1]$ and $A_{1,1} := [b_1, b_2]$. As above, we prove that $h_1(\Omega \setminus A_{1,1}) =]h_1(b_2), h_1(b_1)[\subset A_{1,-1}$ and $h_1^{-1}(\Omega \setminus A_{1,-1}) =]h_1^{-1}(a_1), h_1^{-1}(a_2)[\subset A_{1,1}$. As before, since $\frac{6x^2}{y^2d} < \frac{1}{\rho}$ when $x > 1$, we prove the inclusions.

Since all conditions are satisfied we have, by the Ping-Pong Lemma, that $\langle u, w \rangle$ is a free group. This concludes the proof of the theorem. $\square$

A natural question that can be raised is whether the previous theorem still holds if the norm of the fundamental invertible is -1 . The answer is positive for a 2-Gauss unit with $x \neq 1$. When $\epsilon = 1 + \sqrt{2}$, the same calculations as before can be used to show that $\langle u^2, w \rangle$ is a free group. To see this, we apply the the Ping-Pong Lemma using the following data: $-a_2 = b_2 := 2z_p = 2\sqrt{2}$, $-a_1 = b_1 := \frac{z_0}{2} = \frac{1}{2\sqrt{2}}$, $A_{1,1} := [a_2, a_1]$, $A_{1,-1} := [b_1, b_2]$, $A_{2,1} := [-\infty, a_2[\cup]b_2, \infty]$ and $A_{2,-1} :=]a_1, b_1[$ obtaining that $\langle u^2, w \rangle$ is free. This method does not give us that $\langle u, w \rangle$ is a free group, since it is easily checked, by contradiction, that the only possible choices are $-a_1 = b_1 := \sqrt{2} - 1$ and $-a_2 = b_2 := \sqrt{2} + 1$. However for these values the condition $h_2(a_2) = \varphi_u(a_2) < b_1$ fails to hold.

Corollary 2.4. If $\mathcal{N}(x + y\sqrt{d}) = -1$ and $x \neq 1$ then $\langle u, w \rangle$ is free, where w is the 2-Gauss unit $y\sqrt{-d} + xk$.

PROOF. From $x^2 - y^2d = -1$ we have that $y\sqrt{d} > x$. We apply the proof of the previous theorem using the following data: $-a_2 = b_2 := \frac{3}{2} \frac{y\sqrt{d}}{x}$ and $-a_1 = b_1 := \frac{1}{2} \frac{x}{y\sqrt{d}}$. The condition $x \neq 1$ is equivalent to $\epsilon \neq 1 + \sqrt{2}$. $\square$

We make use of the algebraic properties of the Pell units. This allow us a precise control of their images in $\mathcal{M}$ and to apply the Ping-Pong Lemma in a non-standard way.

Next, we give a criterium for two elements φ_1, φ_2 to generate a free semigroup. Note that we do not use the Ping-Pong Lemma. For an application $\varphi : V \rightarrow V$, we say that $U \subset V$ is invariant under φ if, and only if, the restriction of φ on U is a subset of U .

Lemma 2.5. *Let V be a set of infinite cardinality and let $\varphi_1, \varphi_2 : V \rightarrow V$ be injective maps of infinite order. If $U \subsetneq V$ is invariant under φ_1 and φ_2 , and $x_0 \in V \setminus U$ is a fixed point of φ_1 such that $\varphi_2(x_0) \in U$, then $\langle \varphi_1, \varphi_2 \rangle$ is a free semigroup.*

PROOF. Suppose that the reduced word $\varphi = \varphi_{r_1}^{s_1} \cdots \varphi_{r_k}^{s_k}$ is the identity map. If $r_k = 1$, then, since φ is a reduced word and U is invariant under both maps, $x_0 = \varphi(x_0) = \varphi_{r_1}^{s_1} \cdots \varphi_{r_k}^{s_k}(x_0) = \varphi_{r_1}^{s_1} \cdots \varphi_{r_{k-1}}^{s_{k-1}}(x_0) \in U$. If we have that $r_k = 2$, then $x_0 = \varphi(x_0) \in U$, since $\varphi_2(x_0) \in U$. In any case we have a contradiction because $x_0 \notin U$. $\square$

We now give an application of the above result in the context of the Pell units. Here we do not need to put any restriction on the norm of the fundamental invertible ϵ , unless, like the 4-Pell unit, when the restriction $\mathcal{N}(\epsilon) = 1$ is necessary.

Theorem 2.6. *Let $\epsilon = x + y\sqrt{d}$ be the fundamental invertible of $\mathbb{Q}(\sqrt{d})$ and $u = x + (y\sqrt{-d})i$. If w is one of the units*

- (1.): $y\sqrt{-d} + xk$;
- (2.): $\frac{x+1}{2} - \frac{y\sqrt{-d}}{2}i + \frac{x-1}{2}j + \frac{y\sqrt{-d}}{2}k$;
- (3.): $x^2 - (xy\sqrt{-d})i - y^2dj + (xy\sqrt{-d})k$.

Then $\langle u, w \rangle \subset \mathcal{U}(\frac{-1, -1}{\mathfrak{o}_K})$ is a free semigroup.

PROOF. Consider $\varphi_1, \varphi_2 \in \mathcal{M}, V$ the set of extended real numbers, U the set of positive real numbers and $x_0 = 0$. We claim that the conditions of the previous lemma are satisfied by these data.

In fact, item (1.): $\varphi_1(z) = \varphi_w(z) = \frac{x-y\sqrt{d}}{x+y\sqrt{d}}z$ clearly keeps U invariant and fixes $x_0 = 0 \notin U$. $\varphi_2(z) = \frac{x}{y\sqrt{d}} \frac{\frac{y\sqrt{d}}{x}z+1}{\frac{y\sqrt{d}}{x}z+1}$ has a pole at $-\frac{y\sqrt{d}}{x}$, a zero at $\frac{-x}{y\sqrt{d}}$ and $\varphi_\infty = \frac{y\sqrt{d}}{x}$. If $\mathcal{N}(\epsilon) = 1$, then $\varphi_2(0) = \frac{x}{y\sqrt{d}} > \frac{y\sqrt{d}}{x} > 0$. Hence $\varphi_2(0) \in U$ and $\varphi_2(U \cup \{0\}) =]\frac{y\sqrt{d}}{x}, \frac{x}{y\sqrt{d}}] \subset U$. If $\mathcal{N}(\epsilon) = -1$, then $\frac{y\sqrt{d}}{x} > \frac{x}{y\sqrt{d}} = \varphi_2(0)$. Hence $\varphi_2(0) \in U$ and $\varphi_2(U \cup \{0\}) = [\frac{x}{y\sqrt{d}}, \frac{y\sqrt{d}}{x}[\subset U$.

Item (2.), we set $\varphi_2(z) := \varphi_w^{-1}(z) = \frac{-(y\sqrt{d}-(x+1))z+(y\sqrt{d}-(x-1))}{(y\sqrt{d}+(x-1))z+(y\sqrt{d}+(x+1))}$. Let z_p and z_0 be the pole and the zero of φ_2 , clearly $z_0 < z_p$. Also, $\varphi_\infty = \frac{-(y\sqrt{d}-(x+1))}{(y\sqrt{d}+(x-1))} < \frac{y\sqrt{d}-(x-1)}{y\sqrt{d}+(x+1)} = \varphi_2(0)$. Hence $\varphi_2(U) =]\varphi_\infty, \varphi_2(0)[\subset U$. For the item (3.), define $\varphi_2(z) := \varphi_w^{-1}(z) = \frac{(x^2-xy\sqrt{d})z+(y^2d+xy\sqrt{d})}{(-y^2d+xy\sqrt{d})z+(x^2+xy\sqrt{d})}$. The proof follows as in the item before.

The conditions of the previous lemma being met, it follows that $\langle \varphi_1, \varphi_2 \rangle$ is a free semigroup. $\square$

Let u and v be units in $(\frac{-1, -1}{\mathfrak{o}_{\mathbb{Q}(\sqrt{-2})}})$. It follows from this theorem that if $u = 1 + (\sqrt{-2})i$ and $w = \sqrt{-2} + k$ then the semigroup $\langle u, w \rangle$ is free while we cannot say that these units generate a free group.

References

- [1] Beardon, A. F. *The Geometry of Discrete Groups*, Graduate Texts in Mathematics 91, Springer-Verlag, New York, 1995. xii+337 pp. ISBN: 0-387-90788-2.

- [2] Bridson, M. R., Haefliger, A. *Metric Spaces of Non-Positive Curvature*, Springer, Berlin, 1999.
- [3] Corrales, C., Jespers, E., Leal, G., del Río, Á. *Presentations of the Unit Group of an Order in a Non-Split Quaternion Algebra*, *Advances in Mathematics* 186(2004), 498-524.
- [4] Hartley, B., Pickel, P. F. *Free Subgroups in the Unit Groups of Integral Group Rings*, *Can J. Math.*, vol XXXII, No. 6, 1980, pp 1342-1352.
- [5] Higman, G. *The Units of Group-Rings*, *Proc. London Math. Soc.*, (2)46, (1940), 231-248.
- [6] Juriaans, S. O., Passi, I. B. S., Souza Filho, A. C. *Hyperbolic Unit Groups and Quaternion Algebras*, *Proc. Indian Acad. Sci. (Math. Sci.)*, vol. 119, No. 1, February 2009, pp. 9-22.
- [7] Gonçalves, J. Z., Mandel, A., Shirvani, M. *Free Products Units in Algebras I. Quaternion Algebras*, *J. Algebra* 214 (1999), No. 1, 301 – 316.
- [8] Gonçalves, J. Z., Mandel, A., Shirvani, M. *Free Products Units in Algebras. II. Crossed products*, *J. Algebra* 233 (2000), no. 2, 567 – 593.
- [9] Marciniak, Z. S., Sehgal, S. K. *Constructing Free Subgroups of Integral Group Ring Units*, *Proceedings of the American Mathematical Society* Volume 125, Number 4, April 1997, Pages 1005 – 1009
- [10] Robinson, J. S. Derek, *A course in the theory of groups*, second edition, Springer Verlag, 1995.
- [11] Tits, J. *Free Subgroups in Linear Groups*, *J. Algebra* 20 (1972), pp 250-270.

INSTITUTO DE MATEMÁTICA E ESTATÍSTICA, UNIVERSIDADE DE SÃO PAULO (IME-USP), CAIXA POSTAL 66281,
SÃO PAULO, CEP 05315-970 - BRASIL
E-mail address: `ostanley@ime.usp.br`

ESCOLA DE ARTES, CIÊNCIAS E HUMANIDADES, UNIVERSIDADE DE SÃO PAULO (EACH-USP), RUA ARLINDO
BÉTTIO, 1000, ERMELINDO MATARAZZO, SÃO PAULO, CEP 03828-000 - BRASIL
E-mail address: `acsouzafilho@usp.br`